

This is a repository copy of *Incidental finding of sickle cell trait from an everyday diabetes test – should the family physician report? retest? refer?*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/149643/>

Version: Accepted Version

Article:

Cronin de Chavez, Anna, Dyson, S, Babbington, Fiona et al. (4 more authors) (2019) Incidental finding of sickle cell trait from an everyday diabetes test – should the family physician report? retest? refer? Clinical Diabetes. ISSN 0891-8929

<https://doi.org/10.2337/cd19-0004>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Incidental finding of sickle cell trait from an everyday diabetes test – should the family physician report? retest? refer?

Journal:	<i>Clinical Diabetes</i>
Manuscript ID	CD19-0004.R2
Manuscript Type:	Practical Pointers
Date Submitted by the Author:	n/a
Complete List of Authors:	Cronin de Chavez, Anna; University of Leeds, School of Healthcare, Faculty of Medicine Dyson , Simon; De Montfort University, School of Allied Health Sciences Babbington, Fiona ; Leeds Teaching Hospitals NHS Trust, Blood Sciences Berghs , Maria ; De Montfort University, School of Allied Health Sciences Atkin, Karl; University of York, Health Sciences Whitelaw, Donald; Bradford Teaching Hospitals NHS Foundation Trust, Diabetes and endocrinology Miller , Adrian; Leeds Teaching Hospitals NHS Trust, Blood Sciences

SCHOLARONE™
Manuscripts

Incidental finding of sickle cell trait from an everyday diabetes test – should the family practitioner report? retest? refer?

In Brief

The HbA_{1c} test is increasingly widely used as a diagnostic and screening test for diabetes mellitus type 2 (T2DM) but the presence of haemoglobin variants, such as sickle haemoglobin, can ~~cause some~~ interfere ~~nee~~ with results in some analytical systems. These interferences are ~~sometimes reported~~ ~~on~~ occasionally reported by laborator~~ies~~y results, leading unprepared patients to suspect they ~~are~~ may be sickle cell carriers and seek confirmation through a sickle cell test.

Incidental findings of Hb variants, and the reporting thereof, present multiple ethical challenges to laboratories, medical practitioners, patients and their family members, ~~but and~~ there appears to be no international or national guidelines on how to deal with the reporting of these findings.

This paper explores issues such as whether informed consent is ~~or isn't~~ necessary, ~~;~~ how the results should be communicated, ~~;~~ how the patient may be affected by knowing their carrier status, ~~;~~ the timing of communications, ~~;~~ complications caused by partial results, and being a 'healthy carrier' at the same time as potentially experiencing symptoms.

Introduction

~~The~~ HbA_{1c} ~~test~~ is increasingly widely used as a diagnostic and screening test for T2DM ~~diabetes mellitus type 2~~. With an estimated 8.8% of adults globally having diabetes mellitus, effective screening, diagnosis and monitoring is of major global importance¹. The biomarker of HbA_{1c} refers to glycated haemoglobin A (A_{1c}) molecules and has gained prominence in the diagnosis of T2DM due to certain advantages over plasma glucose testing regimens ~~some advantages over plasma glucose tests~~².

It is well established that that some haemoglobin (Hb) variants, of which there are hundreds, including the clinically relevant HbS, HbE, HbC, and HbD^{3, 4} may interfere with the validity of the HbA_{1c}

results, meaning these ~~testing strategies and tools~~ s should ideally identify variants where they are present ~~necessary~~.

Incidental findings of Hb variants present several ethical challenges for laboratories, medical practitioners, patients and their families ~~members~~. These challenges have, to date, received little attention. This paper discusses some of the advantages of detecting sickle cell trait, identified by the routine ~~triggered through the everyday~~ HbA_{1c} test, but also several ethical dilemmas.

A multidisciplinary team and a patient ~~that was~~ diagnosed as a sickle cell carrier through the HbA_{1c} test worked together to explore these ethical challenges to produce this paper. ~~It and we~~ it is hoped this ~~paper~~ will ~~initiate instigate discussion~~ discourse around the issues presented and ultimately lead to the development of appropriate international guidelines ~~for practice~~.

What does the sickle cell allele have to do with the HbA_{1c} test for type 2 diabetes ~~type 2~~?

Sickle haemoglobin (HbS) is one of the most common haemoglobin variants. Worldwide an estimated 300 million people have sickle cell trait (SCT) and ~4.4 million people have sickle cell disease (SCD), SCD being the overall name for a group of disorders^{5,6}. Sickle cell disease can be life threatening, or increase risk of complications such as stroke, organ failure and acute chest syndrome⁷. Although SCT is usually clinically silent, there are rare ~~complications sequelae e.g. such as~~ haematuria and splenic infarction^{8,9} and ~~potentially~~ a higher risk of T2DM ~~diabetes type 2~~ related complications such as retinopathy, nephropathy, and hypertension¹⁰. The sickle cell allele is present worldwide and w- ~~Whilst~~ there is perhaps more awareness and research of SCT in African American populations, a higher incidence of SCT is also found in Middle Eastern, Mediterranean, Indian and Latin American populations^{6,11}.

Whereas there ~~is a handful of~~ are few methodological variations ~~in~~ of the glucose ~~tolerance test assays~~ used to diagnose diabetes mellitus, the different methods and systems available ~~and vast number of systems used available~~ for the HbA_{1c} test, in comparison, is a challenge in terms of understanding when Hbhaemoglobin variants might interfere with the system in use. According to the National Glycohaemoglobin Standardisation Programme (NGSP) in the U.S.A. there are ~~6-six~~ six method types available for measuring HbA_{1c}: ~~which are~~ immunoassay, borate affinity, enzymatic, ion-exchange high performance liquid chromatography (HPLC) and capillary electrophoresis ~~and-or~~ borate affinity; HPLC. Currently, there are ~~>There are over~~ 200 analytical systems developed by approximately 70 companies, and ~~>150+~~ of the ~~mse systems~~ are available in the USA. There is also a range of susceptibility to interference by, and ability to recognise, Hb variants^{11,12,13,14,15}. Some of these systems are in laboratories, ~~while-and~~ some are used as ~~in~~ point of care situations ~~facilities~~¹⁵⁶. Table 1 below summarises the NGSP's data on HbA_{1c} testing systems:

Table 1. Numbers of methods, systems for HbA_{1c} tests and their interference with variants

Whilst the number of systems with a reported interference by HbS is relatively low, these could be in the more widelycommonly used systems, and the figures demonstrate there is relatively little research to confirm whether or not there is interference of HbS with the majority of the ~~mse systems~~. According to NGSP, only 16% of all systems have been evaluated using robust methods, 57% of these showed an interference with at least one Hbhaemoglobin type and 19% with HbS. Interference was found with more than one method type but the same method came out with no interference with other systems¹⁶⁷. Furthermore, ~~t~~These systems may also require expert interpretation of results. For example, the electrophoretic principle is based on the separation of constituent particles ~~e.g. haemoglobin~~ in a mobile/liquid phase, ~~-and-~~ interaction and subsequent retention ~~by-on~~ a solid phase according to ~~in the system, according to their~~ physicochemical properties. Elution from the solid phase, and subsequent detection, results in a chromatogram where the area of the peak corresponds with concentration of the ~~analytes-compounds~~ detected. ~~E~~The electrophoresis is essentially separating haemoglobins with different properties, such as glycated Hb or variants, which result in different peaks in a

chromatogram. ~~An expert eye~~Expertise is required to ~~properly adequately~~ interpret chromatographs, as some variants may be “hidden” in the HbA peak, ~~potentially-sometimes~~ changing its height, width and/or shape. Therefore it is important for the physician to know if a variant that interferes with the validity of the HbA_{1c} test is present, so that an alternative method or analytical technique may be used.

~~to use an alternative method if screening for diabetes type 2 where there is no variant interference because the results may not be comparable with internationally established reference values based on HbAA studies.~~

Example of an incidental finding of sickle cell trait from a HbA_{1c} test

Against this background of a huge diversity of methods and systems, what might the patient experience be of the result of this test for someone who had **no** prior knowledge of having sickle cell trait (HbAS)? The patient involved in the case described here queried the result of ‘haemoglobin variant detected’ with their family physician, who said they were probably a carrier for sickle cell disease and only on the request of the patient, -referred him for a sickle cell test to confirm this, ~~on request of the patient.~~ On confirmation of SCT status, no ~~support of~~ genetic counselling was offered, ~~to the patient.~~

Below is ~~an example of a~~the chromatogram held in the laboratory files for the above patient, showing a HbS variant detected using the Tosoh (HPLC) system when a HbA_{1c} test was requested by ~~a the~~ primary care practitioner:

Fig 1 Example of chromatogram for a patient with HbAS detected through a HbA_{1c} test, in laboratory files only:-

Fig 2 below shows how the above result was first communicated to the referring practitioner and ultimately the patient who had requested a printout from a series of blood tests.

Fig 2 Extract of lab report showing wording of the HbA_{1c} result sent to the practitioner and then passed onto the patient

The information provided in Fig. 2 on the haemoglobin variant is fairly limited in its usefulness because it does not provide information on what the variant might be, whether there is one copy of the variant allele (heterozygous; e.g. HbAS) or two (homozygous; e.g. HbSS) or a combination of different haemoglobin variants. The information in [Figure 1](#) may be more useful in this respect, but this is not generally released in [laboratory](#) reports due to the expert interpretation often required. Therefore, what primary care practitioners may infer from this may be highly variable, ~~or nothing at all~~, particularly as the result has been classified as 'Normal'. They may choose to ~~or not to~~ discuss further tests to establish which Hb variant the patient has, or the patient may come to their own conclusions with or without the aid of an online search [revealing research, policies and SCT linked death prevention campaigns^{17,18}](#).

There are currently no international guidelines for the reporting of incidental finding of possible [Hbhaemoglobin](#) variants through HbA_{1c} tests. Although the [World Health Organisation](#) ~~W.H.O.~~ guidelines on the use of the HbA_{1c} test refer to interference from haemoglobin variants, they do not advise on if, when or how to inform a primary care practitioner or patient of a variant [that has been detected^{19,9}](#).

Almost a year after the above test ~~was produced, the laboratory~~ ~~he trust~~ involved refined its reporting to practitioners with the following note, where there are heterozygous variants only [e.g.i.e.](#) HbAS. It is highly unlikely that reporting is uniform across hospital [laboratories](#) ~~trusts~~ in the UK even when using the same systems:

Fig 3 Updated wording of the HbA_{1c} result sent to the practitioners a year after the above result

Advantages of detecting a sickle cell trait through the HbA_{1c} test

There could be some advantages for detecting haemoglobin variants through ~~diabetes type 2~~ T2DM screening. The first advantage is that it is important to note for such a patient that a diagnosis of T2DM ~~type 2 diabetes mellitus~~ must not be made using a HbA_{1c} testing system affected by variants. This is important for ~~affected~~ such individuals but also at a population level in regions such as sub-Saharan Africa where between 10% and 40% of people have SCT²⁰²⁰, but many may not know their status as screening programmes are rare. The HbA_{1c} test may still, however, have some use for monitoring a patient who has already been diagnosed with ~~diabetes~~ T2DM because the comparison is then between the patient's own results over time rather than comparing with others.

The HbA_{1c} test is generally not suitable ~~for rapid onset~~ in any clinical conditions ~~situation where there isn't enough HbA with a normal that affects the~~ lifespan (~120 days) of erythrocytes e.g. the presence of ~~such as for~~ some Hb variants, intravascular haemolysis, and in liver disease, or where there is a rapid onset of diabetes ~~but a normal HbA_{1c}~~ such as gestational diabetes and type 1 diabetes (T1DM). The case of HbAS, like that presented here, isn't even as simple as expecting the Hb to be neatly divided into ~~half~~ 50% HbA and ~~50%~~ half HbS, as people with HbAS almost always have more HbA than HbS, and due to in a In a sickle cell genetic carrier HbS levels may vary between 20 and 45%, so there is considerable variation even within those who are HbAS. Interpreting these results requires someone with expertise in haemoglobinopathies^{21,24}. Additionally, most people have a small percentage of foetal haemoglobin (HbF), so a standardised value for HbA_{1c} in HbAS types would probably not be possible. Additionally some point of care HbA_{1c} testing systems, often those used in pharmacies and GP-family physician practices in the UK have no capacity to detect variants, so the *absence* of variant detection should not be assumed to be completely reliable.

The second advantage is that further testing can be done to confirm a carrier status such as SCT. Knowing their carrier status gives ~~them~~ affected individuals the opportunity to make informed reproductive choices and inform family members so they can consider getting tested for SCT. A few countries worldwide have neonatal screening for sickle cell, but these are not always where SCT is most common²²²⁰. Even where newborn screening is offered parents may not have been informed of

the result, understood it or remembered it²²²³. In one report aAn estimated 40% of people in the USA with SCT did not know their status²⁴²³. Therefore, detecting SCT through a relatively common test such as HbA_{1c} could increase the diagnosis and knowledge of carrier status. It could also alert family members of the possibility of a child being born with SCD, especially where newborn screening is absent or sporadic. In sub-Saharan Africa, treatments are limited and an estimated 50–80% of infants born with SCD in Africa die before the age of 5 years of age²⁵²⁴. In 2009 the UN General Assembly declared SCD to be a major public health concern²⁶²⁵ with 1 in 30 deaths of children under-5 worldwide linked to haemoglobin disorders, most of these being SCD²⁷²⁶

Ethical challenges presented by incidental findings of sickle cell trait through the HbA_{1c} test

~~T~~However there may be more challenges than advantages presented by incidental findings of Hb variants, some of which are discussed belowhere.

The first ethical challenge is that any testing in which Hbhaemoglobin variants are incidentally revealed could be considered a genetic test, or a partial one. Some practitioners may not have anticipated this result, gained consent for this kind of genetic screening, or judged consent to be necessary. This may put them in a difficult position as not obtaining consent goes against many guidelines for ~~consent for~~ genetic screening^{2827,28, 29, 30,29,30,31} – whilst leaving the health practitioner unsupported in the absence of any guidelines of what to do when Hb variant ~~results appear~~is present in their patient. This problem, if following such guidelines, –is potentially an additional burden on health care professionals, especially where SCT is most common. Patients are likely to have no idea agreeing for a test for diabetes may throw up an unexpected indication they are a carrier of the sickle cell gene. Therefore patients, as well as practitioners, would have to be fully informed of the consequences of agreeing to such a test. While we note the trait is largely ‘silent’, it does have some

(rare) clinical consequences, which does create the further dilemma of having information which might have a clinical significance, but not passing that information on. Additionally there is a risk of under diagnosis of ~~diabetes type 2~~T2DM with the HbA_{1c} test.³²³¹

The second ethical challenge is that communicating results is not always straightforward. Even within a formalized programme such as the UK's newborn screening programme, concerns are raised about communicating SCT status³³³². Concerns include inconsistencies: ~~variation~~ in who gives the information and how, how much detail to provide, practitioner competence when devolving complex genetic information to non-specialist health professionals³⁴³³, and whether health records of who has SCT are sufficiently robust³⁵³⁴. Although identifying which professional or organisation is best placed to inform the patient of the newly discovered status is important, it is possible that appropriate access to services, such as genetic counselling ~~appropriate for~~ may not actually be available.

A third ethical challenge is that neither the patient nor the practitioner might be able to predict the full social impact of knowing their status as a sickle cell carrier. Subsequent racism and stigma may also be experienced in revealing sickle trait status which the patient may want to avoid³⁶³⁵. Whilst the different clinical impacts of SCT are disputed, there is a clear potential for societal discrimination through carrier status being known in several areas such as employment³⁶³⁷, insurance³⁷³⁸, sports³⁹³⁸, the criminal justice system⁴⁰³⁹ and marriageable status in some communities⁴¹⁴⁰.

A fourth ethical challenge is that patients need to have control over the results and their timing of their release to other family members. The result may perhaps create an unexpected moral dilemma for an individual, about whether to disclose to wider family members or not. For example, the patient in this case was a minor where one parent had requested a printout of a general blood screening where the wording 'haemoglobin variant detected' was included in the 'normal' HbA_{1c} result. A quick online check indicated sickle cell trait to be the most likely explanation for this. Imagine a scenario then where the father of a child he thought was biologically his, received this result and he already knew neither he nor his partner had SCT. This result could have potentially immediate disastrous

consequences for the couple's relationship and family dynamics, even without a second test to confirm the variant type. It could be equally, or perhaps more, devastating for a child to suddenly learn that they are not biologically related. ~~In addition, in the case of a minor both parents may have access to the results, leaving people potentially unprepared for situations such as unexpected paternity.~~ This is one of the reasons why patients and their families need some degree of control over the revealing of carrier status, as well as confirmation of the result on a separate sample (mislabelled samples are common).

A ~~fifth~~another ethical challenge is presented if a patient is not directly told they have a Hb variant, but are told they should not be screened for ~~diabetes type 2-T2DM~~ using the HbA_{1c} test, for whatever reason. With partial information such as this the patient may seek online information and guess they have SCT and anxious that the trait can cause sudden death as reported with some online sources. They may even confuse SCT with SCD and erroneously conclude they have a serious long-term condition. Within this challenge are questions such as: is partial information worse than no information? Can the information be withheld? Should family members be informed?

A ~~sixth~~final ethical challenge could be in the everyday understanding of how SCT affects, or doesn't affect the body. It is confusing for patients when they are told it is both a "diagnosis" and is either mostly benign and/or is a "healthy carrier" state. In addition to numerous research articles available on the internet, of varying quality (based on evidence) and relevance, there are numerous online accounts of SCT patients reporting a range of symptoms they believe to be caused by SCT but not recognised by their physicians. This may leave anyone with SCT worried and unresolved in how they ~~think of~~interpret their ~~body~~ symptoms, and how they interact with family members and their wider community⁴²⁴¹. A call for a resolution to recognition of symptoms was made at a recent symposium including people with the trait: *'You can't use this to discriminate against us, nor will you use it to deny us healthcare'*⁴³⁴².

Ways forward for testing centres, health care providers, policy makers and patients

In conclusion, despite some potential advantages for detecting haemoglobin variants through ~~the~~ HbA_{1c} testing, there are several ethical and practical dilemmas to be navigated. There are questions around what the different HbA_{1c} systems have to offer in terms of diagnosis, if patients should be told, and at what stage, and what counselling and support should they be offered.² As awareness and reporting grows around the various limitations of the HbA_{1c} tests, healthcare services need to plan for the future to ~~ensure-provide~~ patient and practitioner resources and support. Guidelines for screening and reporting of ~~Hbhaemoglobin~~ variants detected in the process of assessing diabetic status currently revealed incidentally in the process of some tests for diabetes type 2 in a more controlled manner urgently need ~~-developmentdrawing-up~~, with the involvement of affected communities, to resolve these problems. This is not an isolated issue, there are other tests that can potentially throw up incidental findings of sickle cell trait, for example where sickled red blood cells were identified in urine and subsequent test revealed SCT⁴⁴³. This article has not covered thalassemias but there is synergy with the situation described here a parallel example in which where a full blood count may be considered an is indirectly ~~a~~ genetic test as it would reveal a patient's status as a thalassaemia carrier. On a much wider scale, with the rise of genomic testing and precision medicine⁴⁵⁴⁴ this kind of predicament with incidental results will only present more ethical, legal and psychosocial problems and would benefit from time and energy being invested in guidance and solutions.

Should therefore the family physician, or the laboratories they work with report Hb variant results? Should they offer a further test to conform which variant they have? Should they refer the patient for genetic counselling?

There appears to be three principle-main choices available in the scenario of the HbA_{1c} test and sickle cell trait. Firstly, either ~~Either:~~ (1) ~~Dispute~~ the very notion of “incidental” genetic findings and

assert that any test that (indirectly) produces genetic findings must be declared to the client as an ~~(indirect)~~ genetic test beforehand and appropriate permission (and counselling?) provided. Secondly, accept ~~Or (2): Accept~~ the concept of “incidental” genetic findings, but then when sickle cell trait confounds the diabetes test, ask the client for permission to investigate possible SCT further with a confirmatory test for ~~sickle-cell trait~~ SCT (and discuss with the client the implications of this) ~~Or (3) Accept~~ Thirdly, accept the concept of “incidental” genetic findings and focus only on the reliability of the information for assessing diabetes, effectively ignoring the sickle cell trait information except for its role as a confounder of the diabetes-relevant information. Option 1 is ethically the purest but potentially expensive, whereas ~~Option 3 seems downright unethical especially where the information could be of potential benefit to the patient and their family, but~~ Option 2 does at least have the possible ethical benefits of ~~affording~~ affording to the client the decision about whether or not to know about a sickle cell trait status ~~(something that newborns aren't asked when sickle-cell trait is an incidental diagnosis to universal newborn screening for sickle-cell disease)~~. Finding ways forward, therefore, would benefit from a dialogue of all parties managing aspects of this test and its results; laboratories, manufacturers, physicians and most importantly people with sickle cell trait, their families and communities.

Key messages

- ~~The~~ HbA_{1c} test is increasingly used as a standard test for ~~diabetes type 2~~ T2DM screening and monitoring and is ~~done performed~~ by a variety of analytical ~~range of~~ methods. However, result validity ~~its validity~~, in some cases, can be affected in the presence of some Hb ~~haemoglobin~~

variants ~~e.g. (such as)~~ sickle haemoglobin (HbS) ~~in some systems~~ and other ~~in~~ conditions where ~~red~~ cell lifespan may be ~~abless than~~ normal, e.g. liver disease.

- Some, but not all, HbA_{1c} test systems detect Hb variants, and further testing may be required to confirm presence of and type of ~~Hbhaemoglobin~~ variants
- There are currently no international, ~~U.S., or U.K. or national level~~ guidelines on how and when incidental findings should be reported to the referring practitioner or patient
- Potential advantages of detecting HbS could be to avoid HbA_{1c} tests in future for ~~diabetes type 2~~ T2DM screening and diagnosis, but the test may still be useful for monitoring HbA_{1c} changes over time within individuals ~~in patients where the diagnosis has already been made~~, and for the detection of SCT and SCD in ~~the wider family group~~ other family members.
- Ethical challenges of detecting HbS include that consent is often not requested; appropriate communication of the result may be difficult and unsupported by services thereafter; unexpected paternity being revealed; accessing incorrect information about SCT symptoms may create anxiety and confusion; being told partial information may lead to incorrect inferences of more serious conditions and revealing sickle trait status may result in a discrimination and stigmatisation by a range of sources and mechanisms.
- Further research is needed to explore these issues for testing centres, practitioners, patients and their family members in more detail and to form a basis for international guidelines.

Acknowledgements

We give our sincere thanks to the individual whose sickle cell trait status was inadvertently discovered as the result of a routine set of blood tests that included the Hb_{A1C} test. Not only have they

been extremely generous in sharing their result and experience with the authors, but they made a significant contribution to the appropriate discussion and presentation of this result in this article~~paper~~. The affected~~is~~ individual is not named as an author for reasons of confidentiality, but would otherwise have been given this status~~included~~. We are also very grateful for the assistance of the laboratory where this test was conducted, which also shall remain nameless due to confidentiality, ~~where this test was conducted~~ and their willingness to be open to sharing the challenges currently faced with such incidental findings. The generosity of this laboratory also extended to inviting the above mentioned individual to an in-depth demonstration of their laboratory procedures from receiving blood samples through to computer analysis.

Author contributions:

A Cronin de Chavez initiated the idea, liaised with the patient mentioned for ~~permissions~~permissions and feedback, wrote much~~a lot~~ of the paper, revisions and carried out the submission.

Atkin K, Babbington F, Berghs M, Dyson SM, Miller A, Whitelaw D all contributed equally to the knowledge required to build the technical information and arguments in this paper. This included working from a basic skeleton version of the problem in focus, and each commenting on and/or re-writing sections of multiple drafts that were prepared before submission. All are in agreement with the submitted version. All participated in answering the editor's and reviewers comments from the first version.

All authors have stated that there is no conflict of interest

Dr. Anna Cronin de Chavez is the guarantor of this work and, as such, had full access to all the patient and laboratory information used in this article and takes responsibility for the integrity of this information

References

- Ogurtsova, K., da Rocha Fernandes, J.D., Huang, Y., Linnenkamp, U., Guariguata, L., Cho, N.H., Cavan, D., Shaw, J.E. and Makaroff, L.E.. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes research and clinical practice* 2017 **128**, pp.40-50.
- ² Bennett, C.M., Guo, M. and Dharmage, S.C.. HbA1c as a screening tool for detection of type 2 diabetes: a systematic review. *Diabetic medicine* 2007 **24**(4), pp.333-343.
- ³ Little, R.R. and Roberts, W.L. A review of variant hemoglobins interfering with hemoglobin A1c measurement. *Journal of Diabetes Science and Technology* 2009. **3**(3), 446 - 451
- ⁴ Nasir, N.M., Thevarajah, M. and Yean, C.Y. Hemoglobin variants detected by hemoglobin A1c (HbA1c) analysis and the effects on HbA1c measurements. *International journal of diabetes in developing countries* 2010, **30**(2), p.86.
- ⁵ GBD 2015 Disease and Injury Incidence and Prevalence, Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study. *The Lancet* 2016 **388** (10053): 1545–1602
- ⁶ Piel, F.B., Patil, A.P., Howes, R.E., Nyangiri, O.A., Gething, P.W., Dewi, M., Temperley, W.H., Williams, T.N., Weatherall, D.J. and Hay, S.I. Global epidemiology of sickle haemoglobin in neonates: a contemporary geostatistical model-based map and population estimates. *The Lancet*, 2013, **381**(9861), pp.142-151.
- ⁷ Ware, R.E., de Montalembert, M., Tshilolo, L. and Abboud, M.R., Sickle cell disease. *The Lancet*, 2017. **390**(10091), 311-323
- ⁸ Naik, R.P. and Derebail, V.K.. The spectrum of sickle hemoglobin-related nephropathy: from sickle cell disease to sickle trait. *Expert review of hematology*, 2017, **10**(12), pp.1087-1094.
- ⁹ Seegars, M.B. and Brett, A.S.. Splenic infarction associated with sickle cell trait at low altitude. *Hematology*, 2015. **20**(10), pp.607-609.

⁰ Skinner, S.C., Diaw, M., Pialoux, V., Mbaye, M.N., Mury, P., Lopez, P., Bousquet, D., Gueye, F., Diedhiou, D., Joly, P. and Renoux, C Increased Prevalence of Type 2 Diabetes–Related Complications in Combined Type 2 Diabetes and Sick Cell Trait. *Diabetes care* 2018 **41**(12) 2595-2602.

¹ ~~Cronin De Chavez, A., Dharni N. The double-edged sword of bloodlines-classification of colour, race and ethnicity in sickle cell research and screening. *The European Journal of Public Health* 2018 **28**(S1): 048-125. 3.11-P3~~

¹¹ Mansour, A. A. "Comparison between two methods of glycosylated hemoglobin estimation among patients with sickle cell trait and diabetes." *J Diabetology* **2** (2013): 3.

¹² National Glycohemoglobin Standardization Program_ Harmonizing haemoglobin A_{1c} testing [2019a](#) <http://www.ngsp.org/interf.asp> Accessed 18th January 2019

³ National Glycohemoglobin Standardization Program Factors that Interfere with HbA_{1c} Test [2019b](#) Results. <http://www.ngsp.org/factors.asp> Accessed 10th June 2019

³⁴ Skinner, S., Pialoux, V., Fromy, B., Sigauo-Roussel, D. and Connes, P., Sickle-cell trait and diagnosis of type 2 diabetes. *The Lancet Diabetes & Endocrinology*. Published online February 16, 2018 [http://dx.doi.org/10.1016/S2213-8587\(18\)30033-0](http://dx.doi.org/10.1016/S2213-8587(18)30033-0)

⁴ ~~NGSP (2019) Factors that Interfere with HbA_{1c} Test Results. <http://www.ngsp.org/factors.asp> Accessed 10th June 2019~~

¹⁵ ~~Mansour, A. A. "Comparison between two methods of glycosylated hemoglobin estimation among patients with sickle cell trait and diabetes." *J Diabetology* **2** (2013): 3.~~

¹⁵⁶ Oxford Health NHS Foundation Trust. 44. Point-of-care HbA_{1c} tests: diagnosis of diabetes. 2016. <https://www.community.healthcare.mic.nihr.ac.uk/reports-and-resources/horizon-scanning-reports/point-of-care-hba1c-tests-diagnosis-of-diabetes> Accessed 28th May 2018

¹⁶⁷ National Glycohemoglobin Standardization Program (2019-) List of NGSP Certified Methods (updated 6/19, listed by date certified) <http://www.ngsp.org/docs/methods.pdf> Accessed 24th May 2019

¹⁴⁸ ~~National Glycohemoglobin Standardization Program (2019) Factors that Interfere with HbA1c Test Results www.ngsp.org/interf.asp~~
~~Accessed Accessed 24th May 2019~~

¹⁷ TS3 Foundation 2019 <https://www.ts3foundation.com/> Accessed 30th July 2019

¹⁸ As One Foundation. Devaughn Darling, 2019
<http://www.asonefoundation.org/about-us/devaughn-darling/> Accessed 30th July 2019

¹⁹⁹ World Health Organisation. Use of Glycated Haemoglobin (HbA1c) in the Diagnosis of Diabetes Mellitus 2011 http://www.who.int/diabetes/publications/report-hba1c_2011.pdf?ua=1

Accessed 22nd November 2017²⁰ Serjeant, GR and Serjeant, BE. Sickle Cell Disease Third Edition Oxford: Oxford University Press.2001

²⁰ ~~Howson CP, Cedergren B, Giugliani R, Huhtinen P, Padilla CD, Palubiak CS, Santos MD, Schwartz IV, Therrell BL, Umemoto A, Wang J. Universal newborn screening: A roadmap for action. Molecular Genetics and Metabolism. 2018 Apr 25.~~

²¹²⁴ Roberts, I. and de Montalembert, M. Sickle cell disease as a paradigm of immigration hematology: new challenges for hematologists in Europe. Haematologica, 2007 **92**(7), pp.865-871.

~~²²²²⁹ Howson CP, Cedergren B, Giugliani R, Huhtinen P, Padilla CD, Palubiak CS, Santos MD, Schwartz IV, Therrell BL, Umemoto A, Wang J. Universal newborn screening: A roadmap for action. Molecular Genetics and Metabolism. 2018 Apr 25.~~

²⁰³ Atkin, K., Ahmad, W.I. and Anionwu, E.N., 1998. Screening and counselling for sickle cell disorders and thalassaemia: the experience of parents and health professionals. Social science & medicine, 1998 **47**(11), pp.1639-1651.

²⁴³ Bleyer, A.J., Reddy, S.V., Sujata, L., Russell, G.B., Akinnifesi, D., Hire, D., Shihabi, Z., Knovich, M.A., Daeihagh, P., Calles, J. and Freedman, B.I., Sickle cell trait and development of microvascular complications in diabetes mellitus. Clinical Journal of the American Society of Nephrology 2010 CJN-08841209.

²⁴⁵ Aygun, B. and Odame, I., 2012. A global perspective on sickle cell disease. Pediatric blood & cancer, **59**(2), pp.386-390.

- ²⁵⁶ United Nations General Assembly. Recognition of sickle-cell anaemia as a public health problem. 2009 Adopted by the General Assembly <http://dag.un.org/handle/11176/172628>
- ²⁶⁷Modell, B. and Darlison, M. Global epidemiology of haemoglobin disorders and derived service indicators. Bulletin of the World Health Organization, 2008. **86**(6), pp.480-487.
- ²⁷ General Medical Council. Consent principles: Patients and Doctors making decisions together 2008
https://www.gmc-uk.org/static/documents/content/Consent_-_English_0617.pdf
- ²⁸ Lappé, M., Gustafson, J.M. and Roblin, R. Ethical and social issues in screening for genetic disease. New England Journal of Medicine, 1972 **286**(21), pp.1129-1132.
- ²⁹⁸ NHS England/PHE. Service specification no.18 NHS Sickle Cell and Thalassaemia Screening Programme 2017 <https://www.england.nhs.uk/wp-content/uploads/2017/06/service-specification-18.pdf> Accessed 26th May 2018
- ²⁹ ~~Lappé, M., Gustafson, J.M. and Roblin, R. Ethical and social issues in screening for genetic disease. New England Journal of Medicine, 1972 **286**(21), pp.1129-1132.~~
- ³⁰ ~~General Medical Council. Consent principles: Patients and Doctors making decisions together 2008~~
~~https://www.gmc-uk.org/static/documents/content/Consent_-_English_0617.pdf~~
- ³¹³⁰ World Health Organisation (2018). Human Genomics in Global Health 2018
<http://www.who.int/genomics/elsi/gentesting/en/> Accessed 26th May 2018
- ³¹³²Lacy, M. E., Wellenius, G. A., Sumner, A. E., Correa, A., Carnethon, M. R., Liem, R. I., ... & Luo, X. Association of sickle cell trait with hemoglobin A1c in African Americans. 2017 JAMA **317** 507-515.
- ³²³³ Chattoo, S., Atkin, K., Dyson, S., Ahmad, W.I. and Anionwu, E.N. Living with Sickle Cell or Beta Thalassaemia Trait: Implications for Identity and Social Life. 2014
- ³³⁴ Kai, J, Ulph, F, Cullinan, T and Qureshi, N. Communication of carrier status information following universal newborn screening for sickle cell disorders and cystic fibrosis: qualitative study of experience and practice. Health Technology Assessment, 2009 **13**(57): 1-82.

- ³⁴⁵ Little, I, Vinogradova, Y, Orton, E, Kai, J and Qureshi, N. Venous thromboembolism in adults screened for sickle cell trait: a population-based cohort study with nested case-control analysis. *BMJ open*, 2017 **7**(3), p.e012665.
- ³⁵⁶ Berghs, M., Dyson, S.M. and Atkin, K. Resignifying the sickle cell gene: Narratives of genetic risk, impairment and repair. *Health*, 2017 **21**(2), pp.171-188.
- ³⁶⁷ Draper, E. *Risky Business: Genetic testing and exclusionary practices in the hazardous workplace* 1991 Cambridge: Cambridge University Press.
- ³⁷⁸ Duster, T. *Backdoor To Eugenics* 2nd Ed. 2003 New York: Routledge
- ³⁸⁹ Carter, B and Dyson, SM. Actor Network Theory, agency and racism: the case of sickle cell trait and US athletics *Social Theory and Health* 2015 **13**(1): 62-77
- ³⁹⁴⁰ Dyson, SM and Boswell, GR. *Sickle Cell and Deaths in Custody* 2009 London: Whiting and Birch
- ⁴⁰⁴¹ Salama, R.A.A. and Saleh, A.K. Effectiveness of premarital screening program for thalassemia and sickle cell disorders in Ras Al Khaimah, United Arab Emirates. *Journal of Genetic Medicine*, 2016 **13**(1), pp.26-30.
- ⁴¹⁴² Dyson SM, Ahmad WIU and Atkin K. Narrative as re-fusion: making sense and value from sickle cell and thalassaemia trait *Health: An Interdisciplinary Journal for the Social Study of Health, Illness and Medicine* 2016 **20**(6): 616-634.
- ⁴²³ Pesante, M. 'Change the narrative'. *Sickle Cell Trait Education Symposium* As One Foundation, Florida State University 2018
- ⁴³⁴ Srivastava, S., Srivastava, R. and Ghorpade, K.G. Sickled erythrocytes in urine as a clue to the diagnosis of sickle cell trait. *Saudi Journal of Kidney Diseases and Transplantation*, 2017 **28**(4), p.909.
- ⁴⁴⁵ Collins, F.S. and Varmus, H. A new initiative on precision medicine. *New England Journal of Medicine*, 2015 **372**(9), pp.793-795.

|

Fig 1 Example of chromatogram for a patient with HbAS detected through a HbA_{1c} test, in laboratory files only




Fig 2 Extract of lab report showing wording of the HbA_{1c} result sent to the practitioner and then passed onto the patient

HbA_{1c} level – IFCC standardized – (KST47) – 34 mmol/mol 20.00-41.00mmol/mol
01 Normal
See comment below
Haemoglobin variant detected. HbA_{1c} may be useful for monitoring diabetic control provided there is no reduction in red cell life. Diabetes is defined by an HbA_{1c} > 47 mmol/mol and optimum glucose control at HbA_{1c} < 59 mmol/mol.
More info go to [www.pathology.\[TRUSTNAME\].nhs.uk](http://www.pathology.[TRUSTNAME].nhs.uk) and search for HbA_{1c}

Fig 3 Updated wording of the HbA_{1c} result sent to the practitioners a year after the above result

Haemoglobin variant detected. Variants have an unpredictable effect on red cell survival and therefore HbA_{1c} cannot be used for diagnosis of diabetes (suggest fasting glucose). However, HbA_{1c} may still be useful for monitoring glycaemic control.

Table 1. Numbers of methods, systems for HbA_{1c} tests and their interference with variants

Number of methods available for HbA _{1c} tests	6 ¹⁷
Number of A _{1c} testing systems manufacturers	Approx. 70 ¹⁷
Number of systems developed to measure A _{1c} (including versions)	Approx. 239 ¹⁷
Number of systems available in the USA (including versions)	Approx. 156 ¹⁷
Number of systems evaluated as reported by NGSP as using ‘robust’ methods	37 (approx. 16% of all systems)
Number of ‘robust’ studies reported by NGSP where there is interference of results by Haemoglobin variants (C, S, E, D, elevated F and Carb)	21 ¹⁸ (57% of systems evaluated)
Number of the above ‘robust’ studies reported evaluating interference of HbS	19 ¹⁸ (19% of systems evaluated)